



PRODUCTS

CATALOGUE

ABOUT US:

Melo Pharmaceuticals Ltd is a registered pharmaceutical company with extensive experience in Human health, Veterinary, Crop health as well as Public health care products.

We import, distribute and sell quality human, animal and crop health care products from qualified manufacturers around the world.

We're the sole distributors of HANVET - Vietnam products in East Africa. We distribute pharmaceutical products, medical sundries and equipment, chemicals, undertake research, as well as agro inputs and farm equipment.

OUR MISSION

To be a pace setting Conglomerate in domestic and International pharmaceutical markets, providing effective and affordable pharmaceuticals products and services to our clients.

OUR VISION

To be a dynamic, Flagship Ugandan Conglomerate providing quality product, services and manufactured goods on domestic and international markets.

ANTIBIOTICS

Antibiotics are medicines that help stop infections caused by bacteria. They do this by killing the bacteria or by keeping them from copying themselves or reproducing.



HANEGG-PLUS

Helps in vitamin, mineral and amino acid deficiencies in birds.

DOSAGE

Preventive dose: 10g/20 liters of drinking water or 10g/10 kg of feed for chickens or ducklings at 2-5 day old and repeated at the age of 4-5 week old.

TYLOSIN-200

For the treatment of:

- **Cattle, goats, sheep:** Complex pneumonia, shipping fever, arthritis, foot-rot; metritis, leptospirosis etc.
- **Pigs:** Bronchopneumonia, enzootic pneumonia, polyarthritis, leptospirosis, pasteurellosis, erysipelas, Vibrio, enterotoxemia etc.
- **Poultry:** Chronic respiratory disease (CRD and complex CRD); sinusitis etc.

DOSAGE

Deep I.M. injection

- **Average:** 1 ml/20 kg b.w, once a day for 3-5 days
- **Cattle:** 3-5 ml/100 kg b.w
- **Pigs, goats:** 2.5 ml/50 kg b.w
- **Piglets, Dogs, Cats:** 2.5 ml/50 kg b.w
- **Poultry:** Dissolve 10ml into 5 liters of drinking water, for 3-5 consecutive days.

Note:

Do not inject more than 10 ml for cattle and 5 ml for pigs in one injection site.



ENROTRIL - 100

- For the treatment and prevention of infections caused by bacteria.
- Complication of infections in respiratory and GI tract in poultry
- Enzootic pneumonia, atrophic rhinitis, diarrhea, septicemia in pigs, calves, kids, lambs.

DOSAGE

Oral administration via drinking water.

- Poultry:

- Average dose is 1 ml/10 kg b.w (10 mg/kg b.w) or 1 ml per 2 liters of drinking water, for 3-5 consecutive days.

- Treatment of colibacillosis or salmonellosis for 5-7 consecutive days. If necessary, the treatment can be repeated after 5 days.
- **Calves, Lambs, Kids:**
- Average dose is 0.5 ml/10 kg b.w for 3 days. Dose may be increased to 5 mg/kg for 5 days for salmonellosis and complicated respiratory disease

- Piglets:

- 0.25 - 0.5 ml/10 kg b.w, for 3-5 days.

- Dogs, Cats:

- 5 mg/kg once daily or a divided dose given twice daily.



WOUNDS & PARASITES SPRAYS



DERMA SPRAY

For the treatment of;

- Skin infections and infected wounds
- Prophylaxis in surgical operations and post-surgical
- Navel disinfection in newborn animals
- Teat sores
- Foot-rot

- Lacerations and abrasions
- After dehorning of calves, deers
- Vesicular diseases
- In cases of bites in poultry flock

DOSAGE

- Shake well before use
- Clean affected area, spray with thin layer formed on the surface, 1-2 times daily, for 3-5 days

FIP-TOX SPRAY

Treatment and control of infestation of fleas, ticks, lice, sarcoptes on dogs, cats.

DOSAGE

- Shake well before use.
- 1ml solution equivalent to 6 pump actuations
- Spray 3-5ml per 1kg b.w depending on the length of the hair.
- Spray the whole body until it is thoroughly wet unless eyes, mouth

- Do not bath before or after spraying 24hours.

Schedule:

When appearance of the parasites.

- **Fleas, ticks:** 2 times, at an interval of 15 Days and upon to the infestation.
- **Lice:** Generally 1 time only
- **Treatment:** 2 times at an interval of 10 days.



DISINFECTANT



HAN-IODINE 10%

- Antiseptic for skin, wounds, membranes pre- and post-operation.
- Navel dipping of newborn animals.
- Lacerations, abrasions, burns, lesions of FMD, foot-rot.
- Prophylaxis of mastitis (teat spray).

- Dermatitis and other skin infections.

DOSAGE

- Shake well before use.
- Topical application, spray onto wounds or lesions, once or 2-3 times daily.

SUPPLEMENTS



BOOST PRO-Probiotics Powder

Helps in;

- Proper Food Digestion and absorption
- Boosting body immunity to diseases
- Fast and proper growth of your animals, birds and fish.



HAN-MILKSTOCK

- Stimulates appetite for the piglets.
- Can work as milk substitute for the piglets in case the sow lacks enough milk.
- For proper and faster growth, gain weight, pink skin, smooth hair, improves body immunity, prevents diarrhea.

Dosage

- **Newborn:** Mix 1 kg of Han-milkstock in 2 liters of clean water, feed pigs at least 5 times / day

- **Pigs during practice:** Mix 1 kg of Hanmilkstock with 15 kg of feed.
- **Post-weaning pig to 70 days old:** Mix 1 kg of Han-milkstock with 25 kg of feed.
- **Poor sows:** Mix 1 kg of Han-milkstock with 30 kg of feed.
- **Newborn calves, goats and sheep:** Mix 1 kg of Han-milkstock with 2.5 liters of clean water.



HAN-PROCARE

- Helps in prevention and treatment of digestive disorders
- Stimulates proper nutrients absorption and reduces harmful gases in the body

HAN-VITALYTE

For the treatment and prevention of:

- Vitamin and mineral deficiencies
- Stress caused for instance by vaccination, transport, re-grouping, high humidity, high temperature or extreme temperature changes, and infectious diseases.

DOSAGE

Oral administration via drinking water for 3-5 days, depended on individual requirements and environmental conditions.

Poultry: 100g per 100 liters of drinking water

Animals: 50-100g per 100 liters of drinking water



SUPPLEMENTS



HAN-ELECTROLYTE

- Electrolyte replacement after extensive loss of body fluids like after diarrhea, enteritis, dehydration, persistent sweating, salivation, vomiting, bleeding...
- Electrolyte imbalance, retarded growth, impaired feed utilization, soft bones, muscle weakness etc.
- Restore electrolyte during stress condition: hot, high humidity, vaccination, regrouping.
- Stabilize acid-base balance in the body in cases of acidosis, imbalance of potassium/sodium ratio.

DOSAGE

Oral administration via drinking water.

- **Poultry:** 100 g/100 liters of drinking water
- **Cattle, pigs:** 100 g/10 liters of drinking water or 50-100 kg b.w



HAN-AMINOVIT

- Improves improper nutritional feed and poor husbandry conditions.
- Improvement of rearing results, feed efficiency, feathering and flesh/body development in poultry, pigs, calves, dairy and cattle
- Vitamin replacement after treatment with anticoccidials or antibiotics.

DOSAGE

Oral administration via drinking water or feed

- **Poultry:** 50-100g per 200 liters of drinking water, for 4-8 consecutive days.
- **Cattle, sheep, goats, pigs:** 100g per 100 liters of drinking water, for 3-5 consecutive days

NOTE:

Prepare fresh solution before use is required. The medicated water should be used within 12 hours.



HAN-GOODWAY

- Contains most important enzymes and probiotics, which improve micro-flora in gut, resulting in better digestion.
- Stimulates proper growth rate, increasing the digestion of protein and hydroxyl-carbon
- Increases milk lactation in sows, cows and goats
- Improves rearing results, feed conversion ratio (FCR)

DOSAGE

Oral administration via drinking water or feed

- **Preventive dose:** 10g per 10 liters of drinking water (equivalent to 1 g/5-10 kg b.w) or 500-700g per 100 kg of feed, daily, throughout rearing period.
- + **Transportation, in hot weather:** use for 3-5 times, weekly.
- + **Diarrhea, infectious diseases, vaccination:** use for 5-7 consecutive days

MELO - PREMIXES



MELO PIGGER PREMIX

Contains a blend of various essential vitamins, minerals, enzymes and toxin binders vital for wholesome pig nourishment and growth

DOSAGE:

- **Weaners:** 1kg per 200kg of pig feeding ratio
- **Growers:** 1kg per 250kg of pig feeding ratio
- **Finishers:** 1kg per 330kg of pig feeding ratio



MELO BROILER PREMIX

It is a pre-mix powder for oral administration through feed.

It contains vitamins and minerals essential for various physiological functions like;

- Optimum growth and production,
- Weight gaining and proper growth promotion



MELO LAYER PREMIX

- Increases egg production
- Promotes the hardness of the egg shell
- Gives an intense color at crocus of the egg
- Prevents cases of rachitis (rickets)
- Increases feeding appetite
- Stimulates metabolism
- Promotes better food absorption in the digestive system

DOSAGE

1 kg per 500 kg of final feed mix



MELO GENERAL PREMIX

High quality general feed premix with all essential amino acids, vitamins and minerals needed to increase productivity in Poultry, Cattle, and Piggyery.

DOSAGE:

2.5kg per tone (1000kg) of feed



MELO PHARMACEUTICALS LTD.

Old Kampala - Plot 31, Martin Road,
Marina Plaza, Basement Floor, Opposite Quality Supermarket,
Tel: 0702 708 714, 0772 822 538
info@melopharmaceuticalsltd.co.ug

Also available at:

EDEN STORES LIMITED

Container Village-Venus Plaza, Basement floor Room 03, below Equity Bank